Date://
RE: Patient Name:
Policy Number / Health Card Number
Re: Request for funding for SpeediCath® catheter, continence supplies
Insurer:
Dear Insurer,
<name> has been assessed as requiring intermittent catheterisation for management of neurogenic bladder. Under <name> current bladder program of <describe current<="" p=""></describe></name></name>
routine>, it has been determined that he/she has experienced the following:
experienced the following.
Recurrent UTI (urinary tract infections) symptoms may include:  • Autonomic Dysreflexia • Fever • Soreness in back or sides • Cloudy urine with increased odour • Onset or increased episodes of incontinence  Catheter related urethral trauma  Difficulty with insertion  Limited dexterity/hand function  Limited transferring function  Spasticity  Other:
SpeediCath is pre-lubricated with a hydrophilic coating and polished eyelets which facilitate ease of insertion and improve patient comfort. Clinical evidence demonstrates a clear difference between single use hydrophilic catheters versus single use uncoated catheters. SpeediCath hydrophilic catheters were shown to reduce risk of urinary track infections in spinal cord injured patients <sup>1,2</sup> . Finally, several studies show reduced risk of urethral trauma by exerting less urethral friction and hence, causing less urethral micro-trauma, irritation and adherence during insertion and withdrawal, measured as withdrawal friction force and haematuria <sup>3,4,5,6,7,8</sup> .
The clinical data also suggests that compared to conventional uncoated catheters, using a SpeediCath catheter may minimize treatment costs, rehabilitation delays and may lower the risk of antibiotic resistance in this patient group.
We therefore request funding for this product (see prescription in the back) for times a day catheterization for a total of catheters per month.
Sincerely,
Doctor:

## References available upon request:

1. Cardenas DD et al. Intermittent Catheterization With a Hydrophillic-Coated Catheter Delays Urinary Tract Infections in Acute Spinal Cord Injury: A Prospective, Randomized, Multicenter Trial. PM&R 2011;3:408-417. 2. De Ridder DJ et al. Intermittent catheterisation with hydrophilic-coated catheters (SpeediCath) reduces the risk of clinical urinary tract infection in spinal cord inured patients: a prospective randomized parallel comparative trial. Eur Urol 2005;48:991-995.

3. Stensballe J et al. Hydrophilic-coated catheters for intermittent catheterization reducComplications of intermittent catheterization: their prevention and treatmente urethral micro trauma: a prospective, randomized, participant-blinded, crossover study of three different types of catheters. Eur Urol 2005;48:978-983.

4. Wyndaele JI. Spinal Cord 2002;40(10):536-541. Complications of intermittent catheterization: their prevention and treatment.

5. Biering-Sørensen F, Nielsen K, Hansen HV. Spinal Cord 1999b;37(4):299-300. Urethral epithelial cells on the surface on hydrophilic catheters after intermittent catheterization: cross-over study with two catheters.

6. Fader M, Moore KN, Cottenden AM, Pettersson L, Brooks R, Malone-Lee J. BJU Int 2001;88(4):373-377. Coated catheters for intermittent catheterization: Smooth or sticky?

7. Bagi P, Hannibalsen J, Permild R, Stilling S, Looms D. Urologia Internationalis. (DOI:10.1159/000321900). Safety of a new compact male intermittent catheter: a randomised, cross-over, single blind study in healthy male

8. Chartier-Kastler E, Lauge I, Ruffion A, Goossens D, Charvier K, Biering-Sørensen F. Spinal Cord 2011, in press. Safety of a new compact catheter for men with neurogenic bladder dysfunction: a randomised, cross-over, open-labelled study.

	Patient Infor	rmation Label	-
R Male		R. canala	
^	. C 12/10 FD (#20422)	Female	
SpeediCath Comp  SpeediCath Comp	act Set 12/18 FR (#28422) act 12/18 FR (#28692)	SpeediCath Compact Se	☐ 12 FR (#28522)
SpeediCath	☐ 10 FR (#27410)		14 FR (#28524)
	12 FR (#27412)	SpeediCath Compact Ev	e 10 FR (#28110) 12 FR (#28112)
	☐ 14 FR (#27414) ☐ 16 FR (#27416)		☐ 14 FR (#28114)
	☐ 18 FR (#27418)	☐ SpeediCath	8 FR (#27508)
SpeediCath Coud			☐ 10 FR (#27510) ☐ 12 FR (#27512)
	☐ 12 FR (#27492) ☐ 14 FR (#27494)		12 FR (#27512) 14 FR (#27514)
	16 FR (#27496)		☐ 16 FR (#27516)
SpeediCath Paedic	atric	☐ SpeediCath Paediatric	6 FR (#27706)
	☐ 8 FR (#27708) ☐ 10 FR (#27710)		□ 8 FR (#27708) □ <sub>10 FR (#27710)</sub>
Other SpeediCath		☐ Other SpeediCath	
Quantity	☐ 1 box ☐ boxes [	☐ Other	
Repeats			
Condition			
☐ Quadripl	egia 🗆 Paraplegia	☐ Multiple sclerosis ☐ Spir	na Bifida
	cancer   Benign prostatic hyperp		ibetes
	cancer 🗆 Other		
Additional info	rmation		

Doctor:

The information in this document is intended for informational purposes only. The provider is ultimately responsible for all information communicated to the payer for reimbursement. As such, the provider should verify coverage criteria and other payer requirements. Coloplast Canada Corporation makes no warranties or guarantees, expressed or implied, concerning the accuracy or appropriateness of this information for any particular use and this information is not intended to provide direction or advice regarding any specific patients' situation. Coverage requirements will vary based on geographic location and reimbursement authority or entity.